



Gastric Tube Insertion

Clinical Indications:

P PARAMEDIC P

- Gastric decompression in intubated patients or for administration of activated charcoal in patients with altered mental status.

Procedure:

1. Estimate insertion length by superimposing the tube over the body from the nose to the stomach.
2. Flex the neck **if not contraindicated** to facilitate esophageal passage.
3. Liberally lubricate the distal end of the tube and pass through the patient's nostril along the floor of the nasal passage. Do not orient the tip upward into the turbinates. This increases the difficulty of the insertion and may cause bleeding.
4. In the setting of an intubated patient or a patient with facial trauma, oral insertion of the tube may be considered or preferred after securing airway.
5. Continue to advance the tube gently until the appropriate distance is reached.
6. Confirm placement by injecting 20cc of air and auscultate for the swish or bubbling of the air over the stomach. Additionally, aspirate gastric contents to confirm proper placement.
7. Secure the tube.
8. Decompress the stomach of air and food either by connecting the tube to suction or manually aspirating with the large catheter tip syringe.
9. Document the procedure, time, and result (success) on/with the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.