

# Pediatric Bradycardia With Poor Perfusion



## History

- Past medical history
- Foreign body exposure
- Respiratory distress or arrest
- Apnea
- Possible toxic or poison exposure
- Congenital disease
- Medication (maternal or infant)

## Signs and Symptoms


- Decreased heart rate
- Delayed capillary refill or cyanosis
- Mottled, cool skin
- Hypotension or arrest
- Altered level of consciousness

## Differential

- Respiratory failure, Foreign body, Secretions, Infection (croup, epiglottitis)
- Hypovolemia (dehydration)
- Congenital heart disease
- Trauma
- Tension pneumothorax
- Hypothermia
- Toxin or medication
- Hypoglycemia
- Acidosis

**Suspected Beta-Blocker or Calcium Channel Blocker**

**Follow Pediatric Toxicology Protocol**



Pediatric Cardiac Protocol Section

# Pediatric Bradycardia With Poor Perfusion



## Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Use Length-based Resuscitation Tape for drug dosages if applicable.**
- **Ensure patent airway, breathing, and circulation as needed. Administer oxygen. Reassess if bradycardia persists after adequate oxygenation and ventilation.**
- **Bradycardia with adequate pulses, perfusion, and respirations requires no emergency intervention. Monitor and continue evaluation with reassessments.**
- **With HR < 60 / min and poor perfusion despite adequate ventilation and oxygenation, begin CPR immediately.**
- **Epinephrine is first drug choice for persistent, symptomatic bradycardia.**
- **Atropine is second choice, unless there is evidence of increased vagal tone or a primary AV conduction block, then given Atropine first.**
- **Transcutaneous pacing:**
  - Indicated if bradycardia is due to complete heart block or other AV blocks which are not responsive to oxygenation, ventilation, chest compressions, or medications. Indicated with known congenital or acquired heart disease.
  - Transcutaneous pacing is not indicated for asystole or bradycardia due to postarrest hypoxic / ischemic myocardial insult or respiratory failure.
  - Pediatric patients requiring external transcutaneous pacing require the use of pads appropriate for pediatric patients per the manufacturers guidelines.
- **Do not delay therapy when bradycardia is evident and no ECG monitor is available.**
- **Vasopressor agents:**
  - Dopamine 5 – 20 mcg / kg / min IV / IO
  - Epinephrine 0.1 – 1 mcg / kg / min IV / IO
  - Norepinephrine 0.1 – 2 mcg / kg / min IV / IO
  - Dose Calculation:  $\text{mL / hour} = \text{kg} \times \text{dose (mcg / kg / min)} \times 60 (\text{min / hr}) / \text{concentration (mcg / mL)}$
- The majority of pediatric arrests are due to airway problems.
- Most maternal medications pass through breast milk to the infant so maintain high-index of suspicion for OD-toxins.
- Hypoglycemia, severe dehydration and narcotic effects may produce bradycardia. Many other agents a child ingests can cause bradycardia, often is a single dose.