



# ADULT AIRWAY

Assess Respiratory Rate, Effort, Oxygenation  
Is Airway/ Breathing Adequate?

YES

Supplemental oxygen  
Goal oxygen saturation  $\geq 92\%$   
Exit to  
Appropriate Protocol(s)

NO

### Basic Maneuvers First

- open airway chin lift/ jaw thrust
- nasal and/ or oral airway(s)
- Bag-valve mask (BVM) +/- PEEP

Spinal Motion Restriction Procedure/ Protocol TB 8  
*if indicated*

Altered Mental Status Protocol UP 4  
*if indicated*

Respiratory Distress with a Tracheostomy Tube  
Protocol AR 10  
*if indicated*

### Capnography Monitoring

- End-tidal (EtCO<sub>2</sub>) monitoring is mandatory following placement of an endotracheal tube.
- EtCO<sub>2</sub> monitoring is mandatory following placement of a BIAD once available on scene.

*Protocols AR 1, 2, and 3 should be utilized together (even if agency is not using Drug Assisted Airway Protocol) as they contain useful information for airway management.*

Airway Obstructed

YES

Airway Foreign Body Obstruction Procedure

A

Direct Laryngoscopy  
*Optional*

P

Airway Cricothyrotomy Surgical Procedure  
*if indicated*

NO

Breathing/ Oxygenation Support needed?

YES

Supplemental Oxygen  
BVM  
Maintain  
Oxygen Saturation  $\geq 92\%$

B

Consider  
Airway NIPPV Procedure  
Airway BIAD Procedure  
*if indicated*

A

Oral/ Nasotracheal Intubation Procedure  
*if indicated*

P

Chest Decompression Procedure  
*if indicated*

Consider  
Airway Drug Assisted Protocol AR 3  
*if available*

Post-intubation/ BIAD Management  
Protocol AR 8  
*if indicated*

Monitor / Reassess  
Supplemental Oxygen  
*if indicated*

NO

Exit to  
appropriate protocol(s)

### Failed Airway:

Unable to Ventilate and Oxygenate  $\geq 90\%$  during or after one (1) or more unsuccessful intubation attempts

and/ or  
Anatomy inconsistent with continued attempts

and/ or  
Three (3) unsuccessful attempts by most experienced Paramedic/ AEMT.

Exit to  
Adult Failed Airway Protocol AR 2

Notify Destination or Contact Medical Control

## AR 1



# ADULT AIRWAY

## Pearls

- See Pearls section of protocols AR 2 and 3.
- For the purposes of this protocol a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of  $\geq 90\%$ , it is acceptable to continue with basic airway measures.
- Ventilation rate should be 10 - 12 per minute to maintain a EtCO<sub>2</sub> of 35 – 45 and avoid hyperventilation.
- **Anticipating the Difficult Airway and Airway Assessment**
  - **Difficult BVM Ventilation (ROMAN):** Radiation treatment/ Restriction; Obese/ Obstruction/ OB – 2d and 3d trimesters/ Obstructive sleep apnea; Mask seal difficulty (hair, secretions, trauma); Age  $\geq 55$ ; No teeth.
  - **Difficult Laryngoscopy (LEON):** Look externally for anatomical problems; Evaluate 3-3-2 (Mouth opening should equal 3 of patients finger's width, mental area to neck should equal 3 of patient's finger's width, base of chin to thyroid prominence should equal 2 of patients finger's width); Obese, obstruction, OB – 2d and 3d trimesters; Neck mobility limited.
  - **Difficulty BIAD (RODS):** Radiation treatment/ Restriction; Obese/ Obstruction/ OB – 2d and 3d trimesters/ Obstructive sleep apnea; Distorted or disrupted airway; Short thyromental distance/ Small mandible.
  - **Difficulty Cricothyrotomy / Surgical Airway (SMART):** Surgery scars; Mass or hematoma, Access or anatomical problems; Radiation treatment to face, neck, or chest; Tumor.
- Complete an Airway Evaluation Form with any BIAD or Intubation procedure where medications are used to facilitate.
- **Nasotracheal intubation:**
  - Procedure requires spontaneous breathing and may require considerable time, exposing patient to critical desaturation.
  - Contraindicated in combative, anatomically disrupted or distorted airways, increased ICP, severe facial trauma, basal skull fracture, and head injury. Orotracheal route is preferred.
- Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.
- If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment).
- AEMT and Paramedics should consider using a BIAD if oral-tracheal intubation is unsuccessful.
- During intubation attempts use External Laryngeal Manipulation to improve view of glottis.
- Gastric tube placement should be considered in all intubated patients if available or time allows.
- It is important to secure the endotracheal tube well to better maintain ETT placement. Manual stabilization of endotracheal tube should be used during all patient moves / transfers.
- **DOPE:** Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.